

Synergizing science and therapy: Neurorepair and neurorehabilitation in spinal cord injury recovery

Jaishriram Rathored¹  , Tanushree Budhbaware¹ , Mithilesh Kumar¹ 

Resumen

Introduction: Devastating motor, sensory, and autonomic deficits with little spontaneous recovery are the outcome of spinal cord injury (SCI). In the past, therapeutic strategies have frequently focused on either biological healing or functional restoration through rehabilitation. Emerging data supports regenerative rehabilitation, which is a synergistic strategy combining regenerative medicine and activity-based rehabilitation, to improve outcomes after spinal cord injury (SCI).

Objective: The purpose of this review is to summarize the most recent research on regenerative rehabilitation techniques for SCI, examining their processes, advancements in translation, and possible applications in clinical settings.

Methods: A comprehensive literature search was conducted on spinal cord injury, regenerative medicine, neurorehabilitation, stem cells, and neuroplasticity, focusing on peer-reviewed publications discussing combination or synergistic therapies.

Results: Stem cell transplantation and rehabilitation training enhance motor recovery, axonal regrowth, synaptic reconfiguration, and task-specific therapy, with neurotrophic factors and bioengineered scaffolds improving endogenous repair.

Conclusion: Regenerative rehabilitation is a paradigm change in the treatment of SCI that combines the reparative potential of regenerative medicine with the plasticity-enhancing benefits of rehabilitation. For such combinatory approaches to continue to be successful, timing, dosage, and individualization must be optimized.

Palabras clave: Spinal cord injury, Regenerative rehabilitation, Neurorepair, Neuroplasticity, Stem cells, Functional recovery.

Sinergia entre ciencia y terapia: neurorreparación y neurorehabilitación en la recuperación de la lesión medular

Abstract

Introducción: los déficits motores, sensoriales y autonómicos devastadores, con escasa recuperación espontánea, son el resultado de la lesión de la médula espinal (LME). En el pasado, las estrategias terapéuticas se han centrado con frecuencia en la curación biológica o en la restauración funcional mediante rehabilitación. Los datos más recientes respaldan la rehabilitación regenerativa, la cual es una estrategia sinérgica que combina la medicina regenerativa y la rehabilitación basada en la actividad, para mejorar los resultados tras la LME.

Objetivo: el propósito de esta revisión es resumir las investigaciones más recientes sobre las técnicas de rehabilitación regenerativa para la LME, examinando sus mecanismos, los avances en la traslación y sus posibles aplicaciones en entornos clínicos.

Métodos: se realizó una búsqueda bibliográfica exhaustiva sobre lesión de médula espinal, medicina regenerativa, neurorrehabilitación, células madre y neuroplasticidad, centrándose en publicaciones revisadas por pares que discutieran terapias combinadas o sinérgicas.

Resultados: el trasplante de células madre y el entrenamiento de rehabilitación mejoran la recuperación motora, el recrecimiento axonal, la reconfiguración sináptica y la terapia específica de tareas, mientras que los factores neurotróficos y los andamios bioingenierizados mejoran la reparación endógena.

Conclusión: la rehabilitación regenerativa representa un cambio de paradigma en el tratamiento de la LME, al combinar el potencial reparador de la medicina regenerativa con los beneficios potenciadores de la plasticidad propios de la rehabilitación. Para que estos enfoques combinados continúen siendo exitosos, deben optimizarse el momento de intervención, la dosificación y la individualización del tratamiento.

Keywords: lesión de médula espinal, rehabilitación regenerativa, neurorreparación, neuroplasticidad, células madre, recuperación funcional.

¹ Central Research Laboratory and Molecular Diagnostics, Datta Meghe Institute of Higher Education and Research, Sawangi (Meghe), Wardha, Maharashtra, India

Correspondencia/Correspondence:

Jaishriram Rathored, Datta Meghe Institute of Higher Education and Research, Sawangi (Meghe), Wardha, Postal code 442001, Maharashtra, India. E-mail: jaishriz@gmail.com

Historia del artículo/ Article info

Received/Recibido: June 27th, 2025
Revised/Revisado: December 10th, 2025
Accepted/Aceptado: January 23d, 2026
Published Online/Publicado: March 5th, 2026

Citation/Citación: Rathored J, Budhbaware T, Kumar M. Synergizing science and therapy: Neurorepair and neurorehabilitation in spinal cord injury recovery.

Acta Neurol Colomb. 2026;42(1):e2016.
<https://doi.org/10.22379/anc.v42i1.1992>



Introduction

One of the most incapacitating neurological disorders is spinal cord injury (SCI), which causes a partial or whole loss of motor, sensory, and autonomic abilities underneath the site of injury. SCI affects between 250,000 and 500,000 people globally each year (1). The combination of rehabilitation and pharmacological or regenerative strategies, termed regenerative rehabilitation, may offer synergistic benefits that surpass the limitations of each approach individually, addressing significant barriers to treatment and rehabilitation with personal and societal implications (2,3).

The goal of neurorepair is to repair the damaged architecture and function of the central nervous system (CNS) via cellular therapies, biomaterials, neurotrophic agents, and genetic interventions. For example, stem cell therapies have demonstrated potential in boosting remyelination, axonal regeneration, and injury environment modification (Figure 1) (4). At the same time, rehabilitation techniques like activity-based therapies, functional electrical stimulation, and locomotor training seek to restore lost functions by utilizing the CNS's plasticity (5). Research indicates that regenerative therapies for severe or chronic spinal cord injury (SCI) are more effective when combined with rehabilitation, due to the principle of activity-dependent plasticity, which enhances the integration and survival of cells in the damaged spinal cord through electrical stimulation or motor training (6).

Additionally, by altering the inflammatory and extracellular matrix environment, rehabilitation may prepare the host tissue to be more responsive to regenerative therapies (7). Task-specific rehabilitation after stem cell transplantation or biomaterial scaffold implantation improves axonal regrowth and functional outcomes in comparison to either treatment alone, as shown by animal models of SCI (8,9). Patients undergoing intense rehabilitation procedures in conjunction with regenerative therapies have demonstrated significant gains in voluntary movement and functional independence in early-stage clinical trials (10). Results emphasize the significance of dosage, timing, and patient-specific factors for improving regenerative rehabilitation. Ongoing challenges include managing injury conditions, identifying optimal treatment times for combined therapies and applying




Figure 1. Neuroregeneration

Source: Adapted from (11).

animal model research to humans. Furthermore, ethical, practical, and regulatory challenges impede the establishment and approval of these rehabilitation methods.

This review evaluates regenerative rehabilitation for spinal cord injury (SCI), emphasizing molecular insights and evidence from preclinical and clinical studies. It presents a unified approach that integrates neurorepair and neurorehabilitation, marking a significant shift towards a collaborative framework for recovery (12).

Methods

A structured literature evaluation of integrative techniques for neurorepair and neurorehabilitation, particularly for spinal cord injury (SCI), was conducted. The methodology that followed involved qualitative synthesis and systematic searches of electronic databases from January 2010 to March 2025, with Boolean operators and MeSH terms being utilized. A snowballing strategy helped identify additional relevant studies, ensuring a coherent thematic synthesis aligned with these research objectives and maintaining scientific rigor.

Criteria for inclusion and exclusion

The eligibility criteria for studies on spinal cord injuries included a focus on regenerative and rehabilitative therapies, unique data from human or animal subjects, and quantifiable results related to neuroplasticity, axon regeneration, motor rehabilitation, or functional enhancement. Studies that were solely pharmaceutical or surgical, lacked detail, or did not provide specific metrics on brain function or repair were excluded.

Data extraction and study selection

After screening the papers by title and abstract, two reviewers conducted full-text reviews, resolving any disagreements by consensus. Data collected from the included studies consisted of the following: SCI model type and severity; regenerative interventions; rehabilitation regimens; primary and secondary outcomes; and key findings. The Cochrane Risk of Bias instrument (RoB 2) was utilized (13).

Evaluation of quality

The SYRCLE risk of bias tool was used in animal research to assess the methodological quality, including elements such as randomization, blinding, and housing conditions (14). The Cochrane Risk of Bias instrument (RoB 2) was used to clinical research (13). To guarantee the validity of the conclusions, only studies with a low or moderate risk of bias were incorporated into the final synthesis.

Data information/synthesis

A qualitative narrative synthesis approach was used to address heterogeneity in research involving various species, intervention types, and outcome measures. Themes were organized based on translational advancement, preclinical discoveries, and synergistic mechanisms, highlighting trends, challenges, and potential directions for future research.

Mechanisms of regenerative rehabilitation's synergistic recovery

Activity-based therapies intended to support functional recovery are combined with cellular or molecular interventions targeted at neuronal repair in regenerative rehabilitation (15). Physical rehabilitation

improves the integration and efficacy of regenerative therapies by encouraging neuronal activity, synaptic remodeling, and axonal sprouting, which is supported by activity-dependent plasticity (3).

By replacing injured cells, promoting axon growth, and modifying the injury microenvironment, neurorepair techniques like stem cell transplantation, growth factor delivery, and biomaterial scaffolds offer the biological substrate for regeneration (4). To direct neuronal reconnection, these treatments frequently call for the right kind of stimulation. When combined with regenerative techniques, rehabilitation interventions such as electrical stimulation, locomotor training, and robotic-assisted movement have been demonstrated to enhance neuronal survival, synaptic strength, and remyelination (Figure 2) (9).

Preclinical research in favour of combined treatments

The advantages of integrating regenerative treatments with rehabilitation have been shown in numerous animal studies. In rats with spinal cord transection, for instance, (8) demonstrated that while neural stem cell transplantation resulted in axonal regeneration, substantial functional recovery was only attained when combined with treadmill training. Similarly, (7) found that when paired with regular physical therapy, bioengineered scaffolds seeded with stem cells improved axonal development and motor outcomes.

Additionally, compared to mice receiving monotherapies, animals receiving combined treatments exhibited increased cortical-spinal tract integrity and greater motor evoked potentials, according to electrophysiological investigations (Figure 3) (17). According to these results, rehabilitation may be crucial for converting biological repair into functional improvements in addition to serving as a supplement to regenerative therapies.

The main non-vascular models of spinal cord injury illustrated involve excitotoxicity from excessive glutamate release, mechanical impact, and compression leading to tissue breakdown. Immunological and inflammatory responses further worsen cellular damage through cytokine signalling. Additionally, the production of free radicals induces oxidative stress, causing further neuronal and glial damage. Apoptosis leads to the progressive loss of these cells, while demyelination disrupts axonal conduction. Together,




Figure 2. A schematic visualization of an animal model's recovery after stroke via regenerative therapy.

Source: Adapted from (16).




Figure 3. Spinal Cord Injury

Source: Adapted from (18).

these mechanisms contribute to functional impairments and hinder neurological recovery after spinal cord injury (Figure 4).

Challenges and clinical translation

Early-phase clinical trials suggest that regenerative rehabilitation may improve voluntary motor control in chronic spinal cord injury (SCI) patients, particularly the use of autologous stem cells alongside locomotor training. However, challenges such as variability in injury severity, timing of intervention, ethical issues with stem cells, and insufficient randomized controlled trials complicate the field. Continued comprehensive research with long-term follow-ups is crucial, as current rodent model findings have limited human relevance. The document also addresses translational challenges and reviews significant clinical trials (19).

Mechanisms of disease initiation and tissue response following neurotrauma

Spinal cord injury (SCI) triggers a complex cascade of secondary damage that includes glutamate ex-

citotoxicity, calcium influx, oxidative stress, mitochondrial dysfunction, and a strong inflammatory response (20,21). Beyond initial mechanical trauma, spinal cord injury leads to further neurological impairments through mechanisms like demyelination, cavitation, and cell death at the lesion site. Neuroinflammation, depicted in Figure 5, involves various cytokines and phenotypic characteristics that contribute to neurodegeneration, which are critical factors to consider during recovery from spinal cord injury.

Astrogliosis: it is characterized by the proliferation of reactive astrocyte, increased levels of glial fibrillary acidic protein (GFAP) and remodelling of the extracellular matrix. It is a key aspect of the tissue response in the myelinated central nervous system (CNS) (22). This because of their function in creating the glial scar and secreting inhibitory chemicals such chondroitin sulfate proteoglycans (CSPGs), reactive astrocytes were formerly thought to be significant inhibitors of regeneration (23). Recent research, however, shows that astrocytes have a dual role: they maintain tissue integrity, stop the spread of inflammation, and promote early repair responses, even though they do limit axonal growth in chronic stages (24,25).




Figure 4. Major non-vascular mechanisms involved in spinal cord injury pathophysiology

Source: Own elaboration.




Figure 5: Neuroinflammation

Source: Own elaboration.

Axonal regeneration: Both internal and external factors restrict the ability to axons to regenerate in the adult central nervous system. Adult neurons have limited regenerative ability by nature, as growth-promoting genes are suppressed. It has been demonstrated that reactivating growth programs is possible with experimental models that target molecular pathways like PTEN/mTOR, SOCS3/STAT3, and KLF family transcription factors (26,27). Axonal extension is hampered by an extrinsic non-permissive environment that is produced by inhibitory signals from myelin debris (such as Nogo-A, MAG, and OMgp) and the glial scar (28,29).

Researches are exploring new approaches to combat these inhibitory effects. These include the enzymatic breakdown of CSPGs using chondroitinase ABC, the administration of neurotrophic factors such as BDNF, NT-3, and GDNF to promote axonal elongation, and the use of biomaterial scaffolds to close the lesion gap (30,31).

In addition to regeneration, axonal plasticity, or the adaptive remodelling of intact neuronal networks, is a crucial component of functional recovery. It has been demonstrated that rehabilitation techniques, including as activity-based therapies and epidu-

ral electrical stimulation, improve synaptic connection and strength in spared pathways (32,33). This neuroplastic adaptation helps to restore motor and sensory functions by compensating for lost connections. In summary, glial reactivity, inflammatory signals, and failure of regeneration interact dynamically in spinal cord injury (SCI). Effective treatments for promoting repair and recovery in patients with neurotrauma require a deep understanding of these mechanisms.

Discussion

Spinal cord injury (SCI) is a severe neurological disorder that causes lasting deficits in motor function, sensory perception, and autonomic control. The absence of standardized, evidence-based rehabilitation protocols, particularly with regard to the timing and sequencing of interventions, has complicated clinical management. Despite physical therapy being widely used in the treatment of SCI, there is no consensus on the optimal starting point for rehabilitation, how to integrate new regenerative techniques, or which patients would benefit most from specific methods (8). Recent advancements in neurorepair, including stem cell transplantation and biomaterial scaffolds,

have increased the range of available treatments. However, these treatments often yield inconsistent functional recovery, highlighting the importance of activity-dependent processes in achieving neurological improvement. Research suggests that regenerative therapies are most effective when combined with timely rehabilitation techniques, resulting in the concept of regenerative rehabilitation (34).

Cellular and molecular processes contribute to the synergy of enhanced synaptic plasticity and improved survival and integration of transplanted cells, due to the activity-dependent release of neurotrophic factors such as brain-derived neurotrophic factor (BDNF) (35). Rehabilitation can alter the inflammatory environment following spinal cord injury. Chronic inflammation can lead to the formation of a glial scar and the production of inhibitory molecules like chondroitin sulfate proteoglycans (CSPGs), which can obstruct axonal regeneration. However, the initial inflammatory response is essential for removing debris (36). Exercise can reduce chronic inflammation, supporting axonal development and repair. The timing of rehabilitation is crucial; early intervention may exacerbate secondary damage, whereas delaying treatment can reduce its effectiveness. Research indicates that the best outcomes with combination therapies can be achieved within a few days to weeks of injury (37).

A review has highlighted the absence of a cohesive model in spinal cord injury (SCI) research that links damage, glial responses and the challenges of regeneration. It proposes a dynamic framework for phase-specific strategies to improve glial regulation, regenerative techniques, and neuroprotection. The findings reveal that astrocytes have dual roles as protective agents and scar formers, which contradicts the previously held view that they have solely inhibitory functions (20,21). Depending on context and timing, inflammation can act as both destructive and reparative. Advances in glial scar remodelling using drugs such as chondroitinase, ABC and intrinsic axonal reprogramming via PTEN/mTOR and KLF signalling, demonstrate the importance of combined, temporally specific therapies (22,23). Rather than total regeneration, neuroplasticity in spared circuits is increasingly recognized as the main driver of functional recovery (24).

Encouraging findings from clinical trials suggest that spinal cord epidural electrical stimulation (EES) can help individuals with chronic spinal cord injury (SCI)

to regain some voluntary motor control, especially when combined with rehabilitative gait training (38). While these treatments do not directly repair the spinal cord, they appear to activate dormant circuitry that may have been stimulated by prior restoration techniques. In a similar vein, (39) showed that prolonged virtual walking could result in a partial neurological recovery for paraplegic patients with the use of BMIs (brain-machine interfaces). These cases demonstrate that combining repair with intensive training can improve the functionality of chronic spinal cord injuries (SCIs), although generalization is hindered by the variability of SCI types. Most biological treatments are still in the experimental stage and produce mixed results. A comprehensive restructuring of clinical practices is essential to merge neurorepair with neurorehabilitation, and advocate for a multidisciplinary strategy. It is vital to manage patient expectations, especially given the rising trend of unregulated stem cell clinics. Innovations such as closed-loop neuromodulation and bioengineered scaffolds, coupled with personalized medicine via biomarkers, could significantly improve rehabilitation outcomes for people with SCI (40).

Alongside electrical stimulation and stem cell therapy, other techniques are crucial for regenerative rehabilitation after spinal cord injury (SCI). Pharmacological treatments such as minocycline, riluzole, and anti-Nogo-A antibodies have shown potential in enhancing neuroprotection and promoting axonal regeneration (41). These substances enhance regenerative outcomes by altering growth-inhibitory, excitotoxic, and inflammatory pathways. New technologies such as brain-computer interfaces (BCIs) and virtual reality (VR)-assisted rehabilitation are being incorporated into the recovery of patients with (SCI) to encourage foster cortical restructuring and motor learning in interactive environments. Future research should explore multimodal approaches that combine biological therapies with cellular, electrical, pharmaceutical, and digital rehabilitation platforms to improve functional recovery for SCI patients (3,39).

Combining neurorepair and neurorehabilitation is essential for effective spinal cord injury (SCI) treatment. Neurorehabilitation addresses functional deficiencies, while neurorepair focuses on structural recovery. Their collaboration is crucial because recovery requires behavioural and biological support, rather than just tissue restoration or movement training (42).

Recent research highlights the importance of combining regenerative and rehabilitative therapies, particularly the use of stem cells alongside rehabilitation techniques such as electrical stimulation and task-specific training. This combined approach has been shown to significantly enhance cell survival, integration, and functional recovery compared to using either treatment alone (9,43). It is believed that activity-induced changes such as the release of neurotrophic factor, synaptic plasticity and neurovascular remodelling contribute to the synergistic effect on recovery. Moreover, the timing and sequencing of interventions, particularly early rehabilitation, have been shown to maximize the benefits of regeneration (44).

Considering the individual diversity in injury level, severity, age, comorbidities, and neuroplastic capability, personalized rehabilitation strategies are essential, highlighting the need for customized approaches for various therapeutic populations (45). For example, due to their greater plasticity, younger patients often respond better to regenerative therapies, whereas older patients may derive greater benefit from assistive technology and medication (46). Recent insights suggest that hormonal factors and healing patterns in spinal cord injury (SCI) outcomes differ by sex. Advanced tools such as wearable sensors, functional imaging and AI evaluations enable real-time monitoring and adaptation of therapy plans to individual needs. Integrating precision medicine techniques into rehabilitation could significantly improve functional recovery and long-term outcomes (47).

It is challenging to find the therapeutic window for integrating regenerative and rehabilitative therapies due to the significant impact of timing on outcomes. For example, early rehabilitation can exacerbate injuries, while delays can reduce neuroplasticity. The absence of standardized protocols complicates clinical trials and reduces the effectiveness of treatment. The proliferation of unregulated stem cell clinics is raising ethical concerns as they are offering untested treatments to vulnerable patients. The new regenerative rehabilitation paradigm emphasizes therapeutic timing, dosage, and individualization, linking rehabilitation directly to neurorepair processes rather than treating it as a mere supportive intervention (48).

In addition to causing immediate suffering through negative consequences, such actions can erode

public confidence in reliable research and delay patients' access to evidence-based care (49). Ethical frameworks emphasize informed consent, patient autonomy, and therapeutic accountability, all of which are often at risk with uncontrolled treatments. To address this issue, increased awareness, regulatory enforcement and patient education are needed, alongside the promotion of rigorous clinical trials and the prevention of misinformation by publications and professional associations (50).

Strengths

This review emphasizes the synergy between neurorepair and neurorehabilitation for spinal cord injury recovery by integrating current research on cellular treatments, biomaterials, and neuroplasticity. It advocates coordinated therapy approaches from a multidisciplinary perspective to enhance functional outcomes.

Limitations

The evaluation process is dominated by preclinical and early-phase clinical studies, which limits applicability to broader patient groups. Variations in damage models, intervention timings and treatment regimens also complicate translation. Additionally, the lack of established protocols for biological and restorative treatments highlights the discrepancy between experimental potential and clinical application.

Future directions

Advances in the treatment of spinal cord injury (SCI) have been made through regenerative rehabilitation, combines neurorepair and neurorehabilitation to improve functional recovery. This approach combines biological repair with activity-based interventions to promote neuroplasticity, and overcome limitations of traditional therapies. Research indicates that regenerative therapies, such as growth factor delivery and stem cell transplantation are most effective in neuronal environments created by rehabilitation. Early intervention can boost neuroplasticity, but challenges such as inflammation and graft rejection in chronic SCI must be managed. This requires strategies to manipulate inhibitory molecules for regeneration (36). Personalization in SCI rehabilitation is complicated by factors like age and comorbidities,

necessitating tailored protocols. Technologies such as wearable sensors, neuroimaging, and biomarker profiling support real-time monitoring and adjustments, enabling customized treatment plans (51). Ethical and regulatory challenges hinder the adoption of stem cell treatments, necessitating standardized manufacturing, long-term safety, and proven efficacy via randomized trials. Rehabilitation research should integrate neurobiological outcomes with traditional measures to substantiate stem cell regeneration mechanisms. Notably, combining electrical neuromodulation with stem cell therapy may yield synergistic effects (52). Targeted and continuous administration of regeneration stimuli is possible by bioengineered scaffolds implanted with genetically modified cells or controlled drug release systems (53).

Emerging technologies like artificial intelligence (AI) and closed-loop systems are significant in rehabilitation. AI analyzes biomechanical data to enhance therapy personalization and predict recovery. Closed-loop neuroprosthetic systems aid in brain plasticity and sensorimotor integration, offering promising outcomes for restoring motor functions in spinal cord injury patients (38,54).

Conclusion

Regenerative rehabilitation combines neurorepair and evidence-based neurorehabilitation to aid recovery from spinal cord injuries (SCIs) by leveraging brain plasticity to promote functional recovery. The review emphasizes the importance of multidisciplinary collaboration between neuroscience, rehabilita-

tion, bioengineering and clinical practice, advocating integrated solutions over isolated approaches. It highlights innovative methods such as neuromodulation and digitally assisted rehabilitation, prioritizing patient-centered outcomes and ethical treatment use. This paradigm is crucial for achieving lasting functional recovery and enhancing the quality of life for SCI patients while adapting therapeutic development to evolving clinical data.

Authors' contributions. Jaishriram Rathore: conceptualization, methodology, illustration, writing original draft, proofreading, and professional guidance throughout the review-writing phase; Tanushree Budhbaware and Mithilesh Kumar contributed to the gathering of data and the creation of this manuscript's pictorial presentations. All authors approve the final draft after it has been reviewed.

Ethical implications. Ethics committee approval was not required for this publication.

Funding. None.

Conflicts of interest. The authors declare that they have no conflict of interest in this study.

AI disclosure statement. The authors declare that no artificial intelligence tools were used in the preparation or writing of this article.

Data availability statement. No data are publicly available. For inquiries regarding any information related to this article, please contact the corresponding author.

References

1. International perspectives on spinal cord injury [Internet]. [cited 2025 May 15]. Available from: <https://www.who.int/publications/i/item/international-perspectives-on-spinal-cord-injury>
2. Behrman AL, Ardolino EM, Harkema SJ. Activity-Based Therapy: From Basic Science to Clinical Application for Recovery After Spinal Cord Injury. *J Neurol Phys Ther JNPT*. 2017 Jul;41 Suppl 3(Suppl 3 IV STEP Spec Iss):S39–45. <https://doi.org/10.1097/NPT.000000000000184>
3. Cramer SC, Sur M, Dobkin BH, O'Brien C, Sanger TD, Trojanowski JQ, et al. Harnessing neuroplasticity for clinical applications. *Brain J Neurol*. 2011 Jun;134(Pt 6):1591–609. <https://doi.org/10.1093/brain/awr039>
4. Assinck P, Duncan GJ, Hilton BJ, Plemel JR, Tetzlaff W. Cell transplantation therapy for spinal cord injury. *Nat Neurosci*. 2017 Apr 25;20(5):637–47. <https://doi.org/10.1038/nn.4541>
5. Martin R, Sadowsky C, Obst K, Meyer B, McDonald J. Functional Electrical Stimulation in Spinal Cord Injury: Top Spinal Cord Inj Rehabil [Internet]. 2012 [cited 2025 May 15];18(1):28–33. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3584753/>. <https://doi.org/10.1310/sci1801-28>

6. Patil N, Korenfeld O, Scalf RN, Lavoie N, Huntemer-Silveira A, Han G, et al. Electrical stimulation affects the differentiation of transplanted regionally specific human spinal neural progenitor cells (sNPCs) after chronic spinal cord injury. *Stem Cell Res Ther* [Internet]. 2023 Dec 20 [cited 2025 May 15];14:378. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10734202/> <https://doi.org/10.1186/s13287-023-03597-w>
7. Silva NA, Sousa N, Reis RL, Salgado AJ. From basics to clinical: a comprehensive review on spinal cord injury. *Prog Neurobiol*. 2014 Mar;114:25–57. <https://doi.org/10.1016/j.pneurobio.2013.11.002>
8. Lu P, Wang Y, Graham L, McHale K, Gao M, Wu D, et al. Long-distance growth and connectivity of neural stem cells after severe spinal cord injury. *Cell*. 2012 Sep 14;150(6):1264–73. <https://doi.org/10.1016/j.cell.2012.08.020>
9. van den Brand R, Heutschi J, Barraud Q, DiGiovanna J, Bartholdi K, Huerlimann M, et al. Restoring voluntary control of locomotion after paralyzing spinal cord injury. *Science*. 2012 Jun 1;336(6085):1182–5. <https://doi.org/10.1126/science.1217416>
10. Curt A, Van Hedel HJA, Klaus D, Dietz V, EM-SCI Study Group. Recovery from a spinal cord injury: significance of compensation, neural plasticity, and repair. *J Neurotrauma*. 2008 Jun;25(6):677–85. <https://doi.org/10.1089/neu.2007.0468>
11. Tam RY, Fuehrmann T, Mitrousis N, Shoichet MS. Regenerative Therapies for Central Nervous System Diseases: a Biomaterials Approach. *Neuropsychopharmacology* [Internet]. 2014 Jan [cited 2025 May 15];39(1):169–88. Available from: <https://www.nature.com/articles/npp2013237> <https://doi.org/10.1038/npp.2013.237>
12. Stewart AN, Gensel JC, Jones L, Fouad K. Challenges in Translating Regenerative Therapies for Spinal Cord Injury. *Top Spinal Cord Inj Rehabil* [Internet]. 2023 [cited 2025 May 15];29(Suppl):23–43. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10759906/> <https://doi.org/10.46292/sci23-000445>
13. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019 Aug 28;366:l4898. <https://doi.org/10.1136/bmj.l4898>
14. Hooijmans CR, Rovers MM, de Vries RBM, Leenaars M, Ritskes-Hoitinga M, Langendam MW. SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol*. 2014 Mar 26;14:43. <https://doi.org/10.1186/1471-2288-14-43>
15. Moritz CT, Ambrosio F. Regenerative Rehabilitation: Combining Stem Cell Therapies and Activity Dependent Stimulation. *Pediatr Phys Ther Off Publ Sect Pediatr Am Phys Ther Assoc* [Internet]. 2017 Jul [cited 2025 May 15];29(Suppl 3 IV STEP 2016 CONFERENCE PROCEEDINGS):S10–5. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5488706/> <https://doi.org/10.1097/PEP.0000000000000378>
16. Ito A, Kubo N, Liang N, Aoyama T, Kuroki H. Regenerative Rehabilitation for Stroke Recovery by Inducing Synergistic Effects of Cell Therapy and Neurorehabilitation on Motor Function: A Narrative Review of Pre-Clinical Studies. *Int J Mol Sci* [Internet]. 2020 Jan [cited 2025 May 15];21(9):3135. Available from: <https://www.mdpi.com/1422-0067/21/9/3135> <https://doi.org/10.3390/ijms21093135>
17. Mancuso R, Santos-Nogueira E, Osta R, Navarro X. Electrophysiological analysis of a murine model of motoneuron disease. *Clin Neurophysiol Off J Int Fed Clin Neurophysiol*. 2011 Aug;122(8):1660–70. <https://doi.org/10.1016/j.clinph.2011.01.045>
18. Tsehay Y, Zeng Y, Weber-Levine C, Awosika T, Kerensky M, Hersh AM, et al. Low-Intensity Pulsed Ultrasound Neuromodulation of a Rodent's Spinal Cord Suppresses Motor Evoked Potentials. *IEEE Trans Biomed Eng* [Internet]. 2023 Jul [cited 2025 May 15];70(7):1992–2001. Available from: <https://ieeexplore.ieee.org/document/10004722> <https://doi.org/10.1109/TBME.2022.3233345>
19. Barker RA, Carpenter MK, Forbes S, Goldman SA, Jamieson C, Murry CE, et al. The Challenges of First-in-Human Stem Cell Clinical Trials: What Does This Mean for Ethics and Institutional Review Boards? *Stem Cell Rep* [Internet]. 2018 May 8 [cited 2025 May 15];10(5):1429–31. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5995446/> <https://doi.org/10.1016/j.stemcr.2018.04.010>
20. Oyibo CA. Secondary injury mechanisms in traumatic spinal cord injury: a nugget of this multiply cascade. *Acta Neurobiol Exp (Warsz)*. 2011;71(2):281–99. <https://doi.org/10.55782/ane-2011-1848>
21. Ahuja CS, Nori S, Tetreault L, Wilson J, Kwon B, Harrop J, et al. Traumatic Spinal Cord Injury—Repair and Regeneration. *Neurosurgery*. 2017 Mar 1;80(3S):S9–22. <https://doi.org/10.1093/neuros/nyw080>
22. Sofroniew MV. Astroglial scar. *Cold Spring Harb Perspect Biol*. 2014 Nov 7;7(2):a020420. <https://doi.org/10.1101/cshperspect.a020420>
23. Silver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci*. 2004 Feb;5(2):146–56. <https://doi.org/10.1038/nrn1326>
24. Anderson MA, Burda JE, Ren Y, Ao Y, O'Shea TM, Kawaguchi R, et al. Astrocyte scar formation aids central nervous system axon regeneration. *Nature*. 2016 Apr 14;532(7598):195–200. <https://doi.org/10.1038/nature17623>
25. Burda JE, Sofroniew MV. Reactive gliosis and the multicellular response to CNS damage and disease. *Neuron*. 2014 Jan 22;81(2):229–48. <https://doi.org/10.1016/j.neuron.2013.12.034>
26. Liu K, Lu Y, Lee JK, Samara R, Willenberg R, Sears-Kraxberger I, et al. PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nat Neurosci*. 2010 Sep;13(9):1075–81. <https://doi.org/10.1038/nn.2603>

27. Sun F, He Z. Neuronal intrinsic barriers for axon regeneration in the adult CNS. *Curr Opin Neurobiol.* 2010 Aug;20(4):510–8. <https://doi.org/10.1016/j.conb.2010.03.013>
28. Filbin MT. Myelin-associated inhibitors of axonal regeneration in the adult mammalian CNS. *Nat Rev Neurosci.* 2003 Sep;4(9):703–13. <https://doi.org/10.1038/nrn1195>
29. Schwab ME. Nogo and axon regeneration. *Curr Opin Neurobiol.* 2004 Feb;14(1):118–24. <https://doi.org/10.1016/j.conb.2004.01.004>
30. Bradbury EJ, Moon LDF, Popat RJ, King VR, Bennett GS, Patel PN, et al. Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature.* 2002 Apr 11;416(6881):636–40. <https://doi.org/10.1038/416636a>
31. Alilain WJ, Horn KP, Hu H, Dick TE, Silver J. Functional regeneration of respiratory pathways after spinal cord injury. *Nature.* 2011 Jul 13;475(7355):196–200. <https://doi.org/10.1038/nature10199>
32. Angeli CA, Boakye M, Morton RA, Vogt J, Benton K, Chen Y, et al. Recovery of Over-Ground Walking after Chronic Motor Complete Spinal Cord Injury. *N Engl J Med.* 2018 Sep 27;379(13):1244–50. <https://doi.org/10.1056/NEJMoa1803588>
33. Edgerton VR, Harkema S. Epidural stimulation of the spinal cord in spinal cord injury: current status and future challenges. *Expert Rev Neurother [Internet].* 2011 Oct [cited 2025 Jun 12];11(10):1351–3. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3361963/> <https://doi.org/10.1586/ern.11.129>
34. Cummings BJ, Uchida N, Tamaki SJ, Salazar DL, Hooshmand M, Summers R, et al. Human neural stem cells differentiate and promote locomotor recovery in spinal cord-injured mice. *Proc Natl Acad Sci U S A.* 2005 Sep 27;102(39):14069–74. <https://doi.org/10.1073/pnas.0507063102>
35. Weishaupt N, Blesch A, Fouad K. BDNF: the career of a multifaceted neurotrophin in spinal cord injury. *Exp Neurol.* 2012 Dec;238(2):254–64. <https://doi.org/10.1016/j.expneurol.2012.09.001>
36. Bradbury EJ, Carter LM. Manipulating the glial scar: chondroitinase ABC as a therapy for spinal cord injury. *Brain Res Bull.* 2011 Mar 10;84(4–5):306–16. <https://doi.org/10.1016/j.brainresbull.2010.06.015>
37. Hutchinson KJ, Gómez-Pinilla F, Crowe MJ, Ying Z, Basso DM. Three exercise paradigms differentially improve sensory recovery after spinal cord contusion in rats. *Brain J Neurol.* 2004 Jun;127(Pt 6):1403–14. <https://doi.org/10.1093/brain/awh160>
38. Wagner FB, Mignardot JB, Le Goff-Mignardot CG, Demesmaeker R, Komi S, Capogrosso M, et al. Targeted neurotechnology restores walking in humans with spinal cord injury. *Nature.* 2018 Nov;563(7729):65–71. <https://doi.org/10.1038/s41586-018-0649-2>
39. Donati ARC, Shokur S, Morya E, Campos DSF, Moiola RC, Gitti CM, et al. Long-Term Training with a Brain-Machine Interface-Based Gait Protocol Induces Partial Neurological Recovery in Paraplegic Patients. *Sci Rep.* 2016 Aug 11;6:30383. <https://doi.org/10.1038/srep30383>
40. Johnson KB, Wei W, Weeraratne D, Frisse ME, Misulis K, Rhee K, et al. Precision Medicine, AI, and the Future of Personalized Health Care. *Clin Transl Sci [Internet].* 2021 Jan [cited 2025 May 15];14(1):86–93. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7877825/> <https://doi.org/10.1111/cts.12884>
41. Casha S, Zygun D, McGowan MD, Bains I, Yong VW, Hurlbert RJ. Results of a phase II placebo-controlled randomized trial of minocycline in acute spinal cord injury. *Brain J Neurol.* 2012 Apr;135(Pt 4):1224–36. <https://doi.org/10.1093/brain/awh072>
42. Lima R, Monteiro A, Salgado AJ, Monteiro S, Silva NA. Pathophysiology and Therapeutic Approaches for Spinal Cord Injury. *Int J Mol Sci [Internet].* 2022 Nov 10 [cited 2025 May 15];23(22):13833. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9698625/> <https://doi.org/10.3390/ijms232213833>
43. Edgerton VR, Tillakaratne NJK, Bigbee AJ, de Leon RD, Roy RR. Plasticity of the spinal neural circuitry after injury. *Annu Rev Neurosci.* 2004;27:145–67. <https://doi.org/10.1146/annurev.neuro.27.070203.144308>
44. Harkema S, Gerasimenko Y, Hodes J, Burdick J, Angeli C, Chen Y, et al. Effect of epidural stimulation of the lumbosacral spinal cord on voluntary movement, standing, and assisted stepping after motor complete paraplegia: a case study. *Lancet Lond Engl.* 2011 Jun 4;377(9781):1938–47. [https://doi.org/10.1016/S0140-6736\(11\)60547-3](https://doi.org/10.1016/S0140-6736(11)60547-3)
45. Behrman AL, Ardolino EM, Harkema SJ. Activity-Based Therapy: From Basic Science to Clinical Application for Recovery After Spinal Cord Injury. *J Neurol Phys Ther JNPT.* 2017 Jul;41 Suppl 3(Suppl 3 IV STEP Spec Iss):S39–45. <https://doi.org/10.1097/NPT.000000000000184>
46. Furlan JC, Kattail D, Fehlings MG. The impact of co-morbidities on age-related differences in mortality after acute traumatic spinal cord injury. *J Neurotrauma.* 2009 Aug;26(8):1361–7. <https://doi.org/10.1177/1545968316644344>
47. Lu DC, Edgerton VR, Modaber M, AuYong N, Morikawa E, Zdonowski S, et al. Engaging Cervical Spinal Cord Networks to Reenable Volitional Control of Hand Function in Tetraplegic Patients. *Neurorehabil Neural Repair.* 2016 Nov;30(10):951–62. <https://doi.org/10.1177/1545968316644344>
48. Turner L. The US Direct-to-Consumer Marketplace for Autologous Stem Cell Interventions. *Perspect Biol Med.* 2018;61(1):7–24. <https://doi.org/10.1353/pbm.2018.0024>

49. Sipp D, Robey PG, Turner L. Clear up this stem-cell mess. *Nature*. 2018 Sep;561(7724):455–7. <https://doi.org/10.1038/d41586-018-06756-9>
50. Hyun I. Allowing innovative stem cell-based therapies outside of clinical trials: ethical and policy challenges. *J Law Med Ethics J Am Soc Law Med Ethics*. 2010;38(2):277–85. <https://doi.org/10.1111/j.1748-720X.2010.00488.x>
51. Kokotilo KJ, Eng JJ, Curt A. Reorganization and Preservation of Motor Control of the Brain in Spinal Cord Injury: A Systematic Review. *J Neurotrauma* [Internet]. 2009 Nov [cited 2025 May 15];26(11):2113–26. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3167869/> <https://doi.org/10.1089/neu.2008.0688>
52. Angeli CA, Boakye M, Morton RA, Vogt J, Benton K, Chen Y, et al. Recovery of Over-Ground Walking after Chronic Motor Complete Spinal Cord Injury. *N Engl J Med*. 2018 Sep 27;379(13):1244–50. <https://doi.org/10.1056/NEJMoa1803588>
53. Tobias IC, Abatti LE, Moorthy SD, Mullany S, Taylor T, Khader N, et al. Transcriptional enhancers: from prediction to functional assessment on a genome-wide scale. *Genome*. 2021 Apr;64(4):426–48. <https://doi.org/10.1139/gen-2020-0104>
54. Capogrosso M, Milekovic T, Borton D, Wagner F, Moraud EM, Mignardot JB, et al. A brain-spine interface alleviating gait deficits after spinal cord injury in primates. *Nature*. 2016 Nov 10;539(7628):284–8. <https://doi.org/10.1038/nature20118>